

Today.

Principle of Induction.(continued.)


$$P(0) \wedge (\forall n \in \mathbb{N})P(n) \implies P(n+1)$$

And we get...

$$(\forall n \in \mathbb{N})P(n).$$

...Yes for 0, and we can conclude Yes for 1...
and we can conclude Yes for 2.....

Climb an infinite ladder?



Your favorite example of forever..or the natural numbers...

$$\begin{array}{c} P(0) \\ \forall k, P(k) \implies P(k+1) \\ P(0) \implies P(1) \implies P(2) \implies P(3) \dots \\ (\forall n \in \mathbb{N})P(n) \end{array}$$

Another Induction Proof.

Theorem: For every $n \in \mathbb{N}$, $n^3 - n$ is divisible by 3. ($3 | (n^3 - n)$).

Proof: By induction.

Base Case: $P(0)$ is " $0^3 - 0$ " is divisible by 3. Yes!

Induction Step: $(\forall k \in \mathbb{N}), P(k) \implies P(k+1)$

Induction Hypothesis: $k^3 - k$ is divisible by 3.

or $k^3 - k = 3q$ for some integer q .

$$\begin{aligned} (k+1)^3 - (k+1) &= k^3 + 3k^2 + 3k + 1 - (k+1) \\ &= k^3 + 3k^2 + 2k \\ &= (k^3 - k) + 3k^2 + 3k \quad \text{Subtract/add } k \text{ (Poll!)} \\ &= 3q + 3(k^2 + k) \quad \text{Induction Hyp. Factor.} \\ &= 3(q + k^2 + k) \quad \text{(Un)Distributive + over } \times \end{aligned}$$

Or $(k+1)^3 - (k+1) = 3(q + k^2 + k)$.

$(q + k^2 + k)$ is integer (closed under addition and multiplication).

$\implies (k+1)^3 - (k+1)$ is divisible by 3.

Thus, $(\forall k \in \mathbb{N})P(k) \implies P(k+1)$

Thus, theorem holds by induction. \square

Four Color Theorem.

Theorem: Any map can be colored so that those regions that share an edge have different colors.



Check Out: "Four corners".

States connected at a point, can have same color.

Quick Test: Which states? Utah. Colorado. New Mexico. Arizona.

Gauss and Induction

Child Gauss: $(\forall n \in \mathbb{N})(\sum_{i=0}^n i = \frac{n(n+1)}{2})$ Proof?

Idea: assume predicate $P(n)$ for $n = k$. $P(k)$ is $\sum_{i=0}^k i = \frac{k(k+1)}{2}$.

Is predicate, $P(n)$ true for $n = k+1$?

$$\sum_{i=0}^{k+1} i = (\sum_{i=1}^k i) + (k+1) = \frac{k(k+1)}{2} + k+1 = \frac{k(k+1)+2(k+1)}{2} = \frac{(k+1)(k+2)}{2}.$$

How about $k+2$. Same argument starting at $k+1$ works!

Induction Step. $P(k) \implies P(k+1)$.

Is this a proof? It shows that we can always move to the next step.

Need to start somewhere. $P(0)$ is $\sum_{i=0}^0 i = \frac{0(0+1)}{2}$ **Base Case.**

Statement is true for $n=0$ ($P(0)$) is true

plus inductive step \implies true for $n=1$ ($P(0) \wedge (P(0) \implies P(1)) \implies P(1)$)

plus inductive step \implies true for $n=2$ ($P(1) \wedge (P(1) \implies P(2)) \implies P(2)$)

...


true for $n=k \implies$ true for $n=k+1$ ($P(k) \wedge (P(k) \implies P(k+1)) \implies P(k+1)$)

...

Predicate, $P(n)$, True for all natural numbers! **Proof by Induction.**

Two color theorem: example.


Any map formed by dividing the plane into regions by drawing straight lines can be properly colored with two colors.



Proper coloring: for each line segment the regions on the two sides have different colors.1

Fact: Swapping red and blue gives another valid colors.

Two color theorem: proof illustration.



Base Case.

1. Add line.
2. Get inherited color for split regions
3. Switch on one side of new line.
(Fixes conflicts along new line, and makes no new ones along previous line.)

Algorithm gives $P(k) \implies P(k+1)$. □

Strengthening Induction Hypothesis.

Theorem: The sum of the first n odd numbers is a perfect square.

Theorem: The sum of the first n odd numbers is n^2 .

k th odd number is $2(k-1)+1$.

Base Case 1 (first odd number) is 1^2 .

Induction Hypothesis Sum of first k odds is perfect square $a^2 = k^2$.



- Induction Step
1. The $(k+1)$ st odd number is $2k+1$.
 2. Sum of the first $k+1$ odds is $a^2 + 2k+1 = k^2 + 2k+1$
????
 3. $k^2 + 2k+1 = (k+1)^2$
... $P(k+1)$!

□

Tiling Cory Hall Courtyard.

Use these L-tiles.

To Tile this 4×4 courtyard.



Can we tile any $2^n \times 2^n$ with L-tiles (with a hole) for every n !

Hole have to be there? Maybe just one?

Theorem: Any tiling of $2^n \times 2^n$ square has to have one hole.

Proof: The remainder of 2^{2n} divided by 3 is 1.

Base case: true for $k=0$. $2^0 = 1$

Ind Hyp: $2^{2k} = 3a+1$ for integer a .

$$\begin{aligned}
 2^{2(k+1)} &= 2^{2k} * 2^2 \\
 &= 4 * 2^{2k} \\
 &= 4 * (3a+1) \\
 &= 12a + 3 + 1 \\
 &= 3(4a+1) + 1
 \end{aligned}$$

a integer $\implies (4a+1)$ is an integer. □

Hole in center?

Theorem: Can tile the $2^n \times 2^n$ square to leave a hole adjacent to the center.


Proof:

Base case: A single tile works fine.

The hole is adjacent to the center of the 2×2 square.

Induction Hypothesis:

Any $2^n \times 2^n$ square can be tiled with a hole at the center.



What to do now???

Hole can be anywhere!

Theorem: Can tile the $2^n \times 2^n$ to leave a hole adjacent *anywhere*.

Better theorem ...better induction hypothesis!


Base case: Sure. A tile is fine. □ □

Flipping the orientation can leave hole anywhere. □ □

Induction Hypothesis:

"Any $2^n \times 2^n$ square can be tiled with a hole **anywhere**."

Consider $2^{n+1} \times 2^{n+1}$ square.



Use induction hypothesis in each.

Use L-tile and ... we are done. □

Strong Induction.

Theorem: Every natural number $n > 1$ can be written as a (possibly trivial) product of primes.

Definition: A prime n has exactly 2 factors 1 and n .

Base Case: $n = 2$.

Induction Step:

$P(n)$ = “ n can be written as a product of primes.”

Either $n + 1$ is a prime or $n + 1 = a \cdot b$ where $1 < a, b < n + 1$.

$P(n)$ says nothing about $a, b!$

Strong Induction Principle: If $P(0)$ and

$$(\forall k \in \mathbb{N})((P(0) \wedge \dots \wedge P(k)) \implies P(k+1)),$$

then $(\forall k \in \mathbb{N})(P(k))$.

$$P(0) \implies P(1) \implies P(2) \implies P(3) \implies \dots$$

Strong induction hypothesis: “ a and b are products of primes”

\implies “ $n + 1 = a \cdot b = (\text{factorization of } a)(\text{factorization of } b)$ ”
 $n + 1$ can be written as the product of the prime factors!

□

Well Ordering Principle and Induction.

If $(\forall n)P(n)$ is not true, then $(\exists n)\neg P(n)$.

Consider smallest m , with $\neg P(m)$, $m \geq 0$

$P(m-1) \implies P(m)$ must be false (assuming $P(0)$ holds.)

This is a proof of the induction principle!

I.e.,

$$(\neg \forall n)P(n) \implies ((\exists n)\neg(P(n-1) \implies P(n))).$$

(Contrapositive of Induction principle (assuming $P(0)$)

It assumes that there is a smallest m where $P(m)$ does not hold.

The Well ordering principle states that for any subset of the natural numbers there is a smallest element.

Examples: even numbers, odd numbers, primes, non-primes, etc..

True for rational numbers? Poll.

Note: can do with different definition of smallest.

For example. Use reduced form: a/b and order by $a + b$.

Well ordering principle.

Thm: All natural numbers are interesting.

0 is interesting...

Let n be the first uninteresting number.

But $n - 1$ is interesting and n is uninteresting,

so this is the first uninteresting number.

But this is interesting.


Thus, there is no smallest uninteresting natural number.

Thus: All natural numbers are interesting.

Tournaments have short cycles

Def: A round robin tournament on n players: every player p plays every other player q , and either $p \rightarrow q$ (p beats q) or $q \rightarrow p$ (q beats p .)

Def: A cycle: a sequence of p_1, \dots, p_k , $p_i \rightarrow p_{i+1}$ and $p_k \rightarrow p_1$.




Theorem: Any tournament that has a cycle has a cycle of length 3.

Tournament has a cycle of length 3 if at all.

Assume the the **smallest cycle** is of length k .

Case 1: Of length 3. Done.

Case 2: Of length larger than 3.



Case 1: “ $p_3 \rightarrow p_1$ ” \implies 3 cycle

Contradiction.

Case 2: “ $p_1 \rightarrow p_3$ ” \implies $k - 1$ length cycle!

Contradicts assumption of smallest $k!$

□

Tournaments have long paths.

Def: A round robin tournament on n players: all pairs p and q play, and either $p \rightarrow q$ (p beats q) or $q \rightarrow p$ (q beats p .)

Def: A Hamiltonian path: a sequence

$$p_1, \dots, p_n, (\forall i, 0 \leq i < n) p_i \rightarrow p_{i+1}.$$



Base: True for two vertices.



(Also for one, but two is more fun as base case!)

Tournament on $n + 1$ people,

Remove arbitrary person \rightarrow yield tournament on $n - 1$ people.

By induction hypothesis: There is a sequence p_1, \dots, p_n

contains all the people where $p_i \rightarrow p_{i+1}$



If p is big winner, put at beginning. Big loser at end.

If neither, find first place i , where p beats p_j .

$p_1, \dots, p_{i-1}, p, p_i, \dots, p_n$ is Hamiltonian path.

□

Horses of the same color...

Theorem: All horses have the same color.

Base Case: $P(1)$ - trivially true.

New Base Case: $P(2)$: there are two horses with same color.

Induction Hypothesis: $P(k)$ - Any k horses have the same color.

Induction step $P(k+1)$?

First k have same color by $P(k)$. 1, 2, 2, 3, ..., $k, k+1$

Second k have same color by $P(k)$. 1, 2, 2, 3, ..., $k, k+1$

A horse in the middle in common! 1, 2, 2, 3, ..., $k, k+1$

All k must have the same color! 1, 2, 3, ..., $k, k+1$

How about $P(1) \Rightarrow P(2)$?

Fix base case.

There are two horses of the same color. ...Still doesn't work!!

(There are two horses is \neq For all two horses!!!)

Of course it doesn't work.

As we will see, it is more subtle to catch errors in proofs of correct theorems!!

Common Knowledge.

Using knowledge about what other people's knowledge (your eye color) is.

On day 1, everyone knows everyone sees more than zero.

On day 2, everyone knows everyone sees more than one.

...

On day 99, everyone knows no one sees 98

since everyone knows everyone else does not see 97...

On day 100, ...uh oh!

Another example:

Emperor's new clothes!

No one knows other people see that he has no clothes.

Until kid points it out.

Sad Islanders...

Island with 100 possibly blue-eyed and green-eyed inhabitants.

Any islander who knows they have green eyes must commit ritual suicide that day.

No islander knows their own eye color, but knows everyone else's.

All islanders have green eyes!

First rule of island: Don't talk about eye color!

Visitor: "I see someone has green eyes."

Result: Poll. On day 100, they all do the ritual.

Why?

Summary: principle of induction.

Today: More induction.

$(P(0) \wedge ((\forall k \in \mathbb{N})(P(k) \Rightarrow P(k+1)))) \Rightarrow (\forall n \in \mathbb{N})(P(n))$

Statement to prove: $P(n)$ for n starting from n_0

Base Case: Prove $P(n_0)$.

Ind. Step: Prove. For all values, $n \geq n_0$, $P(n) \Rightarrow P(n+1)$.

Statement is proven!

Strong Induction:

$(P(0) \wedge ((\forall n \in \mathbb{N})(P(n) \Rightarrow P(n+1)))) \Rightarrow (\forall n \in \mathbb{N})(P(n))$

Also Today: strengthened induction hypothesis.

Strengthen theorem statement.

Sum of first n odds is n^2 .

Hole anywhere.

Not same as strong induction. E.g., used in product of primes proof.

Induction \equiv Recursion.

They know induction.

Thm: If there are n villagers with green eyes they do ritual on day n .

Proof:

Base: $n = 1$. Person with green eyes does ritual on day 1.

Induction hypothesis:

If n people with green eyes, they would do ritual on day n .

Induction step:

On day $n+1$, a green eyed person sees n people with green eyes.

But they didn't do the ritual.

So there must be $n+1$ people with green eyes.

One of them, is me.

Sad.

Wait! Visitor added no information. □

Tiling Cory Hall Courtyard.

